Triclopyr 4 EC

EPA Reg. No. 72167-53-74477

Vegetation Manager Triclopyr 4 EC is an herbicide used to control unwanted woody plants and annual and perennial broadleaf weeds on perennial bluegrass, perennial ryegrass, and tall fescue ornamental turf (including sod farms, commercial turf, and golf courses)

Supplemental Labeling

(Not for use in New York or California)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These new directions can be found on the EPA stamped label dated April 29, 2008.

APPLICATION DIRECTIONS FOR ORNAMENTAL TURF

Refer to Table 2 on the master label for a list of broadleaf weeds controlled by Vegetation Manager Triclopyr 4 EC.

For spot treatments, do not apply more than 2 quarts of Vegetation Manager Triclopyr 4 EC per acre in a single application.

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed 2 or 3 times before being treated. When making applications to mature plants, hard-to-control species, or during drought conditions, use higher rates. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. Do not water for 24 hours after application.

Mixing Instructions

When Vegetation Manager Triclopyr 4 EC is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.
Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of Vegetation Manager Triclopyr 4 EC. Provide moderate agitation while completing the addition of water and during application.

Reseeding Precaution: Do not reseed for 3 weeks after application. (This precaution does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding of 400 lb per acre.)

Broadcast Treatment of Ornamental Turf

Apply ½ to 1 quart per acre of Vegetation Manager Triclopyr 4 EC in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species (see General Use Precautions section of this label) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat- or drought-stress and make repeat applications at least 4 weeks apart.

Tank Mixing: To improve the spectrum of activity, Vegetation Manager Triclopyr 4 EC may be tank mixed at a rate of ½ to 1 pint per acre with recommended rates of low volatile amine or ester formulations of 2,4-D, MCPP, or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Spot Treatment of Ornamental Turf

Mix 3/8 to ¾ ounces of Vegetation Manager Triclopyr 4 EC per 1000 square feet in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. **Note:** Do not apply more than 2 quarts per acre or 1.5 ounces per 1000 square feet of Vegetation Manager Triclopyr 4 EC in a single application.

Vegetation Management, LLC
4515 Falls of Neuse Road, Suite 300, Raleigh NC 27609
T: 919.256.9300 ◆ F: 919.256.9308

Control of Kikuyugrass
Apply Vegetation Manager Triclopyr 4 EC at a rate of ½ to 1 quart per acre. To improve activity, MSMA herbicide may be tank mixed with the ½ quart per acre rate of Vegetation Manager Triclopyr 4 EC. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass
Apply Vegetation Manager Triclopyr 4 EC at the rate of 1 quart per acre. Three to four additional applications at 4 week intervals will be required to give adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate. To improve suppression and control of bermudagrass, 1 quart per acre of Vegetation Manager Triclopyr 4 EC may be tank mixed with a postemergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4 week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf (see **Reseeding Precaution** above).

GENERAL USE PRECAUTIONS (in addition to those found on the label attached to the container)

- Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.
- Vegetation Manager Triclopyr 4 EC may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustine grass, or zoysiagrass, unless turf injury can be tolerated.
- Do not apply Vegetation Manager Triclopyr 4 EC to exposed roots of shallow rooted trees and shrubs.
- Do not apply Vegetation Manager Triclopyr 4 EC to golf course greens.
- Do not apply more than 2 quarts of Vegetation Manager Triclopyr 4 EC per acre in a single application when spot treating.
- Do not make direct applications of Vegetation Manager Triclopyr 4 EC or allow spray mists containing Vegetation Manager Triclopyr 4 EC to drift onto cotton, fruit or orchard trees, shrubs, grapes, peanuts, soybeans, tobacco, vegetable crops, flowers, citrus, or other desirable broadleaf plants.

File: Vegetation Manager Triclopyr 4EC supplemental label for ornamental turf (EPA app 4-29-08)

Vegetation Management, LLC
4515 Falls of Neuse Road, Suite 300, Raleigh NC 27609
T: 919.256.9300 ◆ F: 919.256.9308