

QUALI-PRO™

Quali-Phite® 0-0-26

Foliar Nutrient

FOR NURSERY CROPS, TURF AND COMMERCIAL LANDSCAPING

GUARANTEED ANALYSIS

Soluble Potash (K₂O) 26%
Derived from Potassium Phosphate and Potassium Citrate.

GENERAL RECOMMENDATIONS

Quali-Phite® is a concentrated foliar nutrient solution containing, in highly soluble form Potassium which is beneficial to plant growth, rooting, root development, flowering, transplant establishment and, with sufficient nitrogen, may improve color development when applied as a properly timed foliar or soil spray. Application to turf may improve resistance to traffic if potassium is not adequately supplied by the soil medium. Quali-Phite® is intended for use as a supplemental foliar and/or soil treatment. Incorrect soil application may cause plant injury. Follow Manufacturer's Recommended Directions for Use. Applications of Quali-Phite® should be part of a fertility program based upon soil and tissue testing data. Quali-Phite® may be applied through all types of ground spraying equipment.

Foliar Nutritional Applications - Apply 1 to 4 pints per acre. DO NOT EXCEED a maximum product concentration of 0.1% volume/volume (v/v) for GREENHOUSE or PROTECTED AREA conditions. Note: A 0.1% v/v solution is approximately 0.30 fl. oz. of Quali-Phite® per 2.5 gallons of water. (See APPLICATION PRECAUTIONS Section) DO NOT EXCEED a maximum product concentration of 0.3% volume/volume (v/v) for OUTDOOR or FIELD GROWING conditions. Note: A 0.3% v/v solution is approximately 1 fl. oz. of Quali-Phite® per 2.5 gallons of water. (See APPLICATION PRECAUTIONS Section)

Soil Nutritional Applications - For infrequent applications, DO NOT EXCEED a maximum product concentration of 0.5% volume/volume (v/v). Note: A 0.5% v/v solution is approximately 1.5 fl. oz. of Quali-Phite® per 2.5 gallons of water. (See APPLICATION PRECAUTIONS Section)
For continuous feed application (fertigation), DO NOT EXCEED a maximum product concentration of 0.05% volume/volume (v/v). Note: A 0.05% v/v solution is approximately 0.15 fl. oz. of Quali-Phite® per 2.5 gallons of water. (See APPLICATION PRECAUTIONS Section)

Turf or Tree Farm Nutritional Applications – DO NOT EXCEED a maximum product concentration of 2.0% volume/volume (v/v). Note: A 2.0% v/v solution is approximately 6.5 fl. oz. of Quali-Phite® per 2.5 gallons of water. (See APPLICATION PRECAUTIONS Section)

Compatibility – Quali-Phite® is compatible with most commonly used foliar fungicides, insecticides and chemically neutral fertilizers and microelements. However, the USER ASSUMES FULL RESPONSIBILITY to ensure compatibility when tank mixing with other products. AVOID MIXING with oxidizing chemicals such as chlorine, hydrogen peroxide, ozone and other sterilants. It is not recommended to use this product with surfactants or adjuvants that lower pH. (See APPLICATION PRECAUTIONS Section)

IMPORTANT NOTICE

Seller warrants that this product conforms to its chemical description as stated on the label. The product is reasonably fit for the purpose stated on the label when used in accordance with the label directions and instructions under normal conditions of use. Neither this warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to the seller. Buyer assumes the risk of any such use.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-bw.com>

Quali-Phite® is a registered trademark of Biagro Western Sales, Inc.

US PATENT NO. 5,514,200 and 7,160,349

Manufactured for FarmSaver.com, LLC, 4515 Falls of Neuse Rd., Ste. 300, Raleigh, NC 27609

SUGGESTED USES

CONIFER AND DECIDUOUS WOODY ORNAMENTALS

PROPAGATION – Momentarily submerge cuttings in a 0.3% v/v solution prior to planting.

TRANSPLANTING AND ESTABLISHMENT – Momentarily submerge entire seedling tray in a 0.1% v/v solution OR apply as a foliar spray using a 0.1% v/v solution prior to transplanting.

NEW OUTDOOR PLANTINGS AND RESETS – Once the plants are set, depending on size, apply up to 2 gallons of a 0.5% v/v solution to the root zone of each plant (e.g. 2 pints for small shrubs; 2 gallons for large shrubs). Make additional foliar applications of a 0.1% to 0.3% v/v solution on a ten to fourteen day spray interval or as needed to supplement nutritional requirements.

MAINTENANCE, GROWTH AND DEVELOPMENT – Apply as a foliar spray using a 0.1% to 0.3% v/v solution with subsequent applications at ten to fourteen day intervals or as needed to supplement nutritional requirements.

Note: Applications should be discontinued during dormancy. The first seasonal application should be applied at bud emergence. The last application should be applied just prior to the onset of senescence. For foliar applications, DO NOT EXCEED a product spray concentration of 0.1% to 0.3% v/v depending upon the growing location and/or environment of the plants (See Foliar Nutritional Application Section). When using a 0.3% solution of Quali-Phite®, use a maximum of 4 gallons of spray solution per 1,000 sq. ft.

TREES AND TREE FARMS (e.g. Christmas Trees; Park Plantings)

NEW PLANTINGS AND RESETS – Once the trees are set in the ground, apply up to 2 gallons of a 0.5% v/v solution to the root zone of each tree. Make additional foliar applications of 2 pints per acre at two week spray intervals during the growing season. Make a minimum of two foliar applications per season. Quali-Phite® may be applied as a tank mix partner with foliar fungicide and/or insecticide applications.

MAINTENANCE – Make foliar applications of 2 pints per acre at two week spray intervals during the growing season. Make a minimum of two foliar applications per season. Quali-Phite® may be applied as a tank mix partner with foliar fungicide and/or insecticide applications.

Note: Apply when plants are actively growing. DO NOT EXCEED a maximum product spray concentration of 2.0% v/v. (See Turf or Tree Farm Nutritional Section).

ALL FLOWERING, BEDDING AND NON-WOODY ORNAMENTALS

PROPAGATION – Apply the first foliar spray application at the second true leaf stage using a 0.3% v/v solution. Apply subsequent applications at ten to fourteen day intervals or as needed to supplement nutritional requirements.

TRANSPLANTING AND ESTABLISHMENT – Momentarily submerge the entire seedling tray in a 0.1% v/v solution OR apply as a foliar spray using a 0.1% v/v solution prior to transplanting.

MAINTENANCE, GROWTH AND DEVELOPMENT – Apply as a foliar spray using a 0.1% to 0.3% v/v solution with subsequent applications at ten to fourteen day intervals or as needed to supplement nutritional requirements.

TRANSPORTING – Five to seven days prior to shipment, make a foliar spray application of a 0.1% to 0.3% v/v solution to each plant.

Note: For foliar applications, DO NOT EXCEED a product spray concentration of 0.1% to 0.3% v/v depending upon the growing location and/or environment of the plants (See Foliar Nutritional Application Section). When using a 0.3% solution of Quali-Phite®, use a maximum of 4 gallons of spray solution per 1,000 sq. ft.

CUT FLOWERS

PROPAGATION – Momentarily submerge cuttings in a 0.3% v/v solution prior to planting.

TRANSPLANTING AND ESTABLISHMENT – Momentarily submerge the entire seedling tray in a 0.1% v/v solution OR apply as a foliar spray using a 0.1% v/v solution prior to transplanting.

MAINTENANCE, GROWTH AND DEVELOPMENT – Apply as a foliar spray using a 0.1% to 0.3% v/v solution with subsequent applications at ten to fourteen day intervals or as needed to supplement nutritional requirements.

CUT FLOWER SHIPMENT – Cut flowers may be treated with a 0.1% v/v solution two to three days prior to harvest OR dipped in a 0.1% v/v solution after harvest and prior to shipping.

Note: For foliar applications, DO NOT EXCEED a product spray concentration of 0.1% to 0.3% v/v depending upon the growing location and/or environment of the plants (See Foliar Nutritional Application Section). When using a 0.3% solution of Quali-Phite®, use a maximum of 4 gallons of spray solution per 1,000 sq. ft.

TURF, GOLF COURSE GREENS AND FAIRWAYS

COOL SEASON GRASSES – Apply 3 to 5 fl. oz. per 1,000 square feet in a minimum spray volume of 2 gallons of water at fourteen day intervals during active growth periods.

WARM SEASON GRASSES – Apply 3 to 5 fl. oz. per 1,000 square feet in a minimum spray volume of 2 gallons of water at fourteen day intervals from mid Spring to mid Fall. Continue subsequent applications at 4 to 5 week intervals for remainder of the year.

GOLF GREENS – Depending on the type of grass, follow the suggested use for either COOL or WARM SEASON GRASSES.

TEES AND FAIRWAYS – Apply 3 to 5 fl. oz. per 1,000 square feet in a minimum spray volume of 2 gallons of water during active growth periods.

Note: DO NOT EXCEED a maximum product spray concentration of 2% v/v. (See Turf or Tree Farm Nutritional Section).

IRRIGATION SYSTEM APPLICATIONS

Applications of Quali-Phite® applied through irrigation systems can be substituted for foliar applications described above and can be applied through sprinkler (e.g. solid set, hand move, walking, wheel roll) or low volume (e.g. micro sprinkler, tape, drip, fanjet) irrigation systems. The specific injection equipment and irrigation system must conform to all federal, state and local regulations pertaining to the fertilization of plants through an irrigation system and should be calibrated accurately prior to any application. The irrigation system should be operated to achieve maximum distribution uniformity to avoid plant injury and/or ineffectiveness. For best results, apply Quali-Phite® toward the end of the irrigation cycle allowing sufficient time for the product to be flushed out of the entire irrigation system before turning off the irrigation water.

APPLICATION PRECAUTIONS

Due to the continuous development of new varieties of each of the many different ornamental plants, it is recommended that prior to the first use on a particular ornamental variety, the grower evaluate plant response(s) by treating a small sample of plants (6 to 10) with the product at the recommended dose(s). Allow sufficient time after application of the product to determine if there is going to be any undesirable plant response(s), including but not limited to stunting and/or leaf burn.

AS WITH ALL ACID MATERIALS, USER SHOULD EXERCISE PRECAUTION TO AVOID POTENTIAL FOLIAR BURNING. PRECAUTION IS ADVISED IF APPLYING THIS PRODUCT IN COMBINATION WITH OTHER MATERIALS SINCE MIXING COULD RESULT IN A POTENTIALLY HAZARDOUS OR UNDESIRABLE REACTION CAUSED BY PRODUCT INCOMPATIBILITIES. A COMPATIBILITY TEST IS RECOMMENDED BEFORE TANK MIXING WITH OTHER PRODUCTS. THE pH OF ANY FINAL SPRAY SOLUTION SHOULD BE CHECKED AND AN APPROPRIATE AMOUNT OF THE SOLUTION BE APPLIED TO A SMALL AREA OF THE INTENDED CROP TO ENSURE PLANT COMPATIBILITY PRIOR TO THE PLANNED APPLICATION. USER ASSUMES FULL RESPONSIBILITY TO ENSURE PRODUCT, EQUIPMENT AND PLANT COMPATIBILITIES.

CAUTION! May be irritating to contaminated skin and eyes. Inhalation of sprays or mists may be harmful. Wear gloves and goggles. Do not get in eyes, on skin, or on clothing. Avoid breathing splash, sprays, or mists generated from this material. Store in tightly closed container. Wash thoroughly after handling. If spilled on skin or splashed in eyes, rinse affected area for at least 15 minutes. TARGET ORGANS: Skin and eyes.

Use in well ventilated area. If involved in a fire, this material may decompose and produce irritating fumes and toxic gases (phosphorus oxides, carbon monoxide, and carbon dioxide). In the event of a spill, clean-up promptly using trained personnel and pre-planned procedures or contact CHEMTREC at 1-800-424-9300 for assistance. Consult Material Safety Data Sheet for further information. KEEP OUT OF REACH OF CHILDREN. This Product Intended For Agricultural Use Only.